

## Alterations of Mental Status and Thyroid Hormones after Thermal Injury\*

GEORGE M. VAUGHAN, ARTHUR D. MASON, JR., WILLIAM F. McMANUS, AND  
BASIL A. PRUITT, JR.

U.S. Army Institute of Surgical Research, Brooke Army Medical Center, Fort Sam Houston, Texas 78234

**ABSTRACT.** In 16 burn patients, mean values for serum  $T_4$  and  $T_3$ , their  $T_3$  uptake-derived free indices ( $FT_4I$  and  $FT_3I$ ) and dialysis-derived free concentrations ( $FT_4$  and  $FT_3$ ) were depressed (all  $P < 0.001$ ) compared to respective means in 13 normal subjects. In the patients, the free hormone indices were relatively more depressed below control values than were the free hormone concentrations. However, within the group of burn patients, variation in  $FT_4I$  reflected that of  $FT_4$  ( $r = 0.91$ ), and variation in  $FT_3I$  reflected that of  $FT_3$  ( $r = 0.93$ ). We then studied serum  $T_4$ ,  $T_3$ , and their free indices in 134 patients (burn size, 6–96% of the skin area), including 45 nonsurvivors, none of whom received steroid, dopamine, or iodine treatment. At each sampling, the level of obtundation (LO) was determined on a 6-point scale from normal to deep coma. Whereas initially low

mean  $FT_4I$  values rose in survivors, they remained lower in nonsurvivors than in survivors until death in the nonsurvivors. In nonsurvivors, mean LO worsened in the first week and remained worse than that in survivors until death. Multiple regression analyses showed that for a given age or burn size, nonsurvival was better correlated with lower  $T_4$  or  $FT_4I$  than with  $T_3$  or  $FT_3I$ , but was even more closely correlated with worse LO ( $P < 0.001$ ). Exclusion of data obtained within 24 h of narcotic or tranquilizer doses did not weaken the relationship of nonsurvival with LO and  $FT_4I$ . Nonsurvival after burn injury was associated with reduced  $T_4$ ,  $FT_4I$ , and mental status for up to weeks before death, this association being independent of treatment with drugs acting on mental status or thyroid function. (*J Clin Endocrinol Metab* 60: 1221, 1985)

$T_4$  AND  $T_3$  concentrations in serum may be depressed in burn patients (1–3), similar to thyroid hormones in patients with many other types of nonthyroidal illness (NTI) (4, 5). Previous studies of a variety of NTI patients (6–8) suggested that low serum  $T_4$  concentrations were particularly related to nonsurvival. We sought to determine whether mental status is related to serum thyroid hormone values in burn patients and whether these variables are related to outcome.

### Subjects and Methods

Morning serum samples were obtained approximately once per week for thyroid hormone and other routine chemical determinations in patients in the intensive care areas of our burn unit. These patients were treated with fluids and electrolytes for initial resuscitation and subsequent evaporative loss and were provided with vigorous nutritional support (2), alternating topical mafenide acetate and silver sulfadiazine, systemic antibiotics if needed for infection, morphine (occasionally

meperidine) as needed for pain, and antacids and occasionally cimetidine to prevent Curling's ulcers, and excision and grafting of wounds were performed when indicated. Treatment did not include topical or systemic agents containing iodine. The data reported herein are from 134 adult burn patients without head injury or a history of thyroid disease and with serum TSH levels less than  $20 \mu\text{U/ml}$ , from whom there were 656 samples taken without prior therapy with dopamine, corticosteroids, or other agents to support the cardiovascular system. Forty-five patients were nonsurvivors and died from 5–104 days after injury. Samples were also taken from 13 normal laboratory personnel.

The level of obtundation (LO) in the patients was determined at the time of sampling according to the system used by Rudman *et al.* (9) for patients with head trauma (Table 1). Serum  $T_4$ ,  $T_3$ ,  $T_3$  charcoal uptake ( $T_3U$ ), TSH (Diagnostic Products, Los Angeles, CA), and  $rT_3$  (Serono, Milan, Italy) were determined in serum by RIA (radioassay for  $T_3U$ ) with kits from the manufacturers. Indices of free  $T_4$  and  $T_3$  ( $FT_4I$  and  $FT_3I$ , respectively) were the product of the serum  $T_4$  or  $T_3$  level and  $T_3U$  divided by the  $T_3U$  value of the normal reference sample in the kit.  $T_3U$  was calculated as the tracer counts bound to the matrix divided by the total counts.

In the control and some of the burn samples, the dialyzable fractions of  $T_4$  and  $T_3$  after equilibrium at 37°C were determined at the Nichols Institute (San Juan Capistrano, CA). Free hormone concentrations ( $FT_4$  and  $FT_3$ ) were calculated as the product of the respective hormone concentration and the di-

Received September 29, 1984.

Address requests for reprints to: Dr. George M. Vaughan, Library Branch, U.S. Army Institute of Surgical Research, Fort Sam Houston, Texas 78234-6200.

\* The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. Investigators adhered to AR 70-25 and USAMRDC Reg 70-25 on Use of Volunteers in Research, as well as the Declaration of Helsinki.

TABLE 1. Determination of mental status

Criteria	Level of obtundation
Oriented to person and place, cooperative, not hallucinating (all true)	
no	yes.....0
Conscious when undisturbed	
no	yes.....1
Temporary arousal on minor disturbance	
no	yes.....2
Semipurposeful movement on noxious stimuli	
no	yes.....3
Deep tendon reflexes present	
no	yes.....4
	.....5

alyzable fraction. These samples were assessed with analysis of covariance to determine the correlations of FT<sub>4</sub>I with FT<sub>4</sub> and FT<sub>3</sub>I with FT<sub>3</sub> and assess any influence of burn injury on these relationships. The interrelationships among the measured variables within the larger group of burn patients (excluding data from nonburned subjects) were determined by stepwise multiple regression analysis (10). Survival was coded as 0, and death was coded as 1. In some cases, *t* tests were used to compare two independent means.

### Results

Table 2 and Fig. 1 represent only the samples in which values for dialyzable fractions of T<sub>4</sub> and T<sub>3</sub> were available. Concentrations of T<sub>4</sub> and T<sub>3</sub> were depressed in burn patients, and the T<sub>3</sub>U and dialyzable fractions were elevated. Nevertheless, the free indices (FT<sub>4</sub>I and FT<sub>3</sub>I) and free concentrations (FT<sub>4</sub> and FT<sub>3</sub>) in the patients were also highly significantly depressed compared with normal values. Although in Table 2, the degree of depression of FT<sub>4</sub> appears almost equivalent to that of FT<sub>4</sub>I, analysis of covariance indicates that in the patients, the FT<sub>4</sub>I was relatively more depressed than was the FT<sub>4</sub> level (Fig. 1). The same relationships were evident between FT<sub>3</sub>I and FT<sub>3</sub>. Because it has been suggested that use of T<sub>3</sub>U expressed as the matrix to serum ratio of tracer counts sometimes produces a better correlation between free indices and free concentrations than does

TABLE 2. Results in normal subjects and those burn patients in whom samples for equilibrium dialysis determinations were available

	Unburned normal (n = 13)	Burn survivors (n = 13)	Burn nonsurvivors (n = 3)
T <sub>4</sub> (μg/dl)	7.59 ± 0.35	3.52 ± 0.26 <sup>a</sup>	1.84 ± 0.37
FT <sub>4</sub> I	6.79 ± 0.25	3.56 ± 0.29 <sup>a</sup>	1.90 ± 0.27
FT <sub>4</sub> (ng/dl)	2.14 ± 0.11	1.25 ± 0.08 <sup>a</sup>	0.77 ± 0.13
T <sub>3</sub> (ng/dl)	141 ± 7.2	60 ± 7.2 <sup>a</sup>	32 ± 9.0
FT <sub>3</sub> I	126 ± 5.2	60 ± 7.4 <sup>a</sup>	34 ± 8.6
FT <sub>3</sub> (pg/dl)	316 ± 12	200 ± 19 <sup>a</sup>	146 ± 41
T <sub>3</sub> U (%)	27.1 ± 0.7	30.7 ± 0.8 <sup>b</sup>	33.1 ± 2.4
T <sub>4</sub> DF (%)	0.0282 ± 0.0006	0.0361 ± 0.0007 <sup>a</sup>	0.0440 ± 0.0021
T <sub>3</sub> DF (%)	0.227 ± 0.008	0.347 ± 0.012 <sup>a</sup>	0.471 ± 0.050
Age (yr)	34.1 ± 2.0	35.7 ± 5.3	66.3 ± 13.5
TBS (%)		34.4 ± 4.1	48.3 ± 9.2
PBD		13.5 ± 3.0	4.3 ± 2.0

Values are the mean ± SE. T<sub>4</sub>DF, T<sub>4</sub> dialyzable fraction; T<sub>3</sub>DF, T<sub>3</sub> dialyzable fraction. See text for definition of other variables. In five patients in whom two values for some variables were available, the mean of the two values was used.

<sup>a</sup> *P* < 0.001 vs. normal.

<sup>b</sup> *P* < 0.01 vs. normal.

T<sub>3</sub>U expressed as the matrix to total ratio (4), the above results were reanalyzed employing T<sub>3</sub>U expressed as the matrix to serum ratio. Since the results were the same, including the disproportionately greater depression of free indices relative to the free concentrations in burns, the free indices reported herein are based on the T<sub>3</sub>U as the matrix to total ratio.

Figure 2 shows the hormone and LO values from the first 30 postburn days for patients in the full study group, arranged according to total burn size (TBS; as percentage of body surface area) and survival. The patterns for T<sub>4</sub> and T<sub>3</sub> depression in the burn patients were quite similar to those for FT<sub>4</sub>I and FT<sub>3</sub>I, respectively. Excepting values in the largest burn size group, mean T<sub>4</sub>, T<sub>3</sub>, FT<sub>4</sub>I, and FT<sub>3</sub>I (and LO in each group) were significantly different in nonsurvivors compared with values in survivors (*P* < 0.01). Figure 3 shows FT<sub>4</sub>I and LO (restricted to patients with TBS of 30–60%) according to postburn day (PBD) in the first 3 weeks. Whereas FT<sub>4</sub>I was lowest initially and rose in survivors during the first week, it remained low in nonsurvivors. LO worsened in nonsurvivors during the first week, and survivors had much less obtundation throughout. Both FT<sub>4</sub>I and LO were altered in nonsurvivors for weeks before death and exhibited no additional deterioration in the weeks or days before death. Multiple regression analyses detected no relationship between days before death and FT<sub>4</sub>I or LO. TSH values (mean ± SE; microunits per ml) were not significantly different between survivor (3.26 ± 0.12) and nonsurvivor (3.03 ± 0.30) samples (normal reference range, 0–10). All patients had TSH values less than 20, a practical limit below which TSH has been considered compatible with




FIG. 1. Covariance analysis of free indices (FT<sub>4</sub>I and FT<sub>3</sub>I) and respective free concentrations (FT<sub>4</sub> and FT<sub>3</sub>) of T<sub>4</sub> and T<sub>3</sub>. Within each panel, the slopes did not differ between normal subjects and burn patients, so the lines have a common slope. The positional differences were significant, as indicated. Normal reference ranges: FT<sub>4</sub>I, 5.5-14; FT<sub>4</sub>, 1.32-3.8 ng/dl; FT<sub>3</sub>I, 94-169; and FT<sub>3</sub>, 230-660 pg/dl.

NTI (5).

Multiple regression analyses indicated that FT<sub>4</sub>I was most strongly negatively (–) correlated with LO and that the residual variation in FT<sub>4</sub>I was explained by further correlation with PBD (+), TBS (–), nonsurvival (–), and age (–) (all  $P < 0.001$ ). LO correlated best with nonsurvival (+) and additionally with age (+), PBD (+), FT<sub>4</sub>I (–), and FT<sub>3</sub>I (–) (all  $P < 0.001$ ). Nonsurvival was most strongly correlated with LO (+) and additionally with burn size (+) and FT<sub>4</sub>I (–) (all  $P < 0.001$ ). Other analyses were performed, excluding data collected within 24 h of administration of narcotics or tranquilizers, leaving 183 of the total 656 samples for analysis. FT<sub>4</sub>I was lower with greater burn size, LO was higher in the nonsurvivors, and nonsurvival related separately to greater obtundation and lower FT<sub>4</sub>I (all  $P < 0.001$ ). Cimetidine was administered within 24 h before collec-


tion of 84 of the total 656 samples. After elimination of these 84 samples, multiple regression analyses of the remaining data still indicated correlation (all  $P < 0.001$ ) of FT<sub>4</sub>I with TBS and LO, of LO with survival, and of survival with LO and FT<sub>4</sub>I.

### Discussion

Depression of serum T<sub>3</sub> (1, 2, 3, 11) and T<sub>4</sub> (2, 3) and their free indices in burn patients has been reported previously. We confirm these findings, which are similar to those in other forms of NTI (4, 5). Further, we now report depression of mean serum FT<sub>4</sub> in burns, a finding similar to that reported in other forms of severe illness (12-14), though FT<sub>4</sub> may not always be depressed in NTI (4). Perhaps because of the possibility that circulating inhibitors of thyronine binding to circulating transport proteins in patients with NTI (15, 16) may inhibit binding to the matrix in the T<sub>3</sub>U test (17), the FT<sub>4</sub>I may exhibit a disproportionately greater reduction than that of the dialysis-based FT<sub>4</sub> concentration in these patients. Despite this discrepancy, patients with a wide variety of critical NTI and reduced serum T<sub>4</sub> concentration reportedly had mean reductions not only of FT<sub>4</sub>I but also of FT<sub>4</sub> (12-14). Thus, burn injury is similar to other severe NTI not only in causing reduction of mean total T<sub>3</sub> and T<sub>4</sub> concentrations, but also with respect to the presence of a defect in binding of thyronines in the circulation (elevated dialyzable fractions of T<sub>4</sub> and T<sub>3</sub>), somewhat disproportionate lowering of free indices more than free concentrations, and a depressed mean FT<sub>4</sub> and FT<sub>3</sub>. The previously reported absence of an increased concentration of TSH and its response to TRH or a frankly depressed TSH response to TRH in burn patients represents a further similarity with NTI (3). Burn injury may represent a good model of severe NTI.

Burn patients had depressed FT<sub>4</sub> and FT<sub>3</sub> levels and even more depressed FT<sub>4</sub>I and FT<sub>3</sub>I values as a group compared to values in normal subjects. However, there were proportional variations in FT<sub>4</sub>I with FT<sub>4</sub> and in FT<sub>3</sub>I with FT<sub>3</sub> in the burn patients tested and within a separate larger group of patients (3) not included in the present report. It is clear that among burn patients, those with lower FT<sub>4</sub>I values also have lower FT<sub>4</sub> levels compared to FT<sub>4</sub> levels in those with higher FT<sub>4</sub>I values, and the same applies to FT<sub>3</sub>I and FT<sub>3</sub>. Thus, variations in the free hormone indices within the overall group of burn patients in this study was considered to reflect variations in respective free concentrations. Among the hormones, nonsurvival correlated best with depression of FT<sub>4</sub>I, suggesting lower FT<sub>4</sub> in nonsurviving compared to surviving burn patients. Among survivors of burns, lower T<sub>4</sub>, FT<sub>4</sub>I, T<sub>3</sub>, and FT<sub>3</sub>I were proportional to burn size, with values for nonsurvivors much lower at any given burn size below

FIG. 2. Mean  $\pm$  SE for measured variables during the first 30 PBDs in patients grouped according to burn size and outcome. In patients with more than 1 value for a variable, the mean value was used. The numbers near the symbols represent the number of patients.  $\bullet$ , Survivors;  $\circ$ , nonsurvivors. The values above the abscissa are the mean number of days before death for sampling times in the nonsurvivors. The mean value for 13 normal subjects (CON) is shown (—), as is the extent of 2 SD from the normal mean (---).



60% or time after injury beyond the second day. Since all sampling was restricted to times when the patients were in the intensive care area, there was selection for patients who were more critically ill even among the survivors. This makes the  $FT_4I$  difference between nonsurvivors and survivors more noteworthy and probably explains the apparently blunted rise of  $FT_4I$  after the first week in the survivors.

In reports of other NTI (6, 18), low serum  $T_4$  levels, sampled within 2 days of admission to intensive care units, occurred mainly in those patients who ultimately died. Kaptein *et al.* (7) found that in patients with various NTI admitted to intensive care units with low  $T_4$  levels,  $T_4$  returned toward normal in survivors, but not in nonsurvivors, and the patients' lowest  $T_4$  value correlated negatively with mortality. Our finding of greater reduction of  $T_4$  and  $FT_4I$  in nonsurviving burn patients than in survivors, manifested throughout their course, further indicates similarity between burn injury and other NTI. Calvano *et al.* (8) also found lower  $T_4$  and  $FT_4I$  in nonsurviving than in surviving burn patients.

In patients with head trauma, serum  $T_4$ ,  $FT_4I$ , and  $T_3$  were reduced in proportion to the degree of coma (9). It is interesting to note the reported difference in patients with coma due to head trauma exhibiting an elevated TSH response to TRH (9) compared to burn patients exhibiting a normal (or in nonsurvivors, a frankly blunted) TSH response (3). It is possible that in burns and other NTI, the metabolic milieu somehow suppresses the normal ability of thyrotrophs to recognize a deficiency of thyroid hormone and consequently augment their response to TRH. Perhaps this ability is not lost in patients whose pathological process may be a lesion restricted to the head and causing both coma and relative deficiency of endogenous hypothalamic TRH.

Reductions of mental status and of circulating  $T_4$  can also be associated in the burn model of NTI. Greater reduction of  $FT_4I$  and mental status were both associated with mortality. Of course, one explanation of our results might be that these associations merely stem from otherwise unrelated parallel changes in  $FT_4I$ ,  $LO$ , and mortality accompanying greater severity of injury. However,




FIG. 3. Mean  $\pm$  SE of FT<sub>4</sub>I and LO for patients with TBS of 30-60% grouped by outcome and PBD during the first 3 weeks postinjury. The numbers by the symbols are the number of patients in each group. The numbers above the abscissa are the mean number of days before death (DBD) at which the samples were taken in nonsurvivors. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$  (by  $t$  test; survivors vs. nonsurvivors).

regression analysis indicated variation in survival associated with variation in mental status and FT<sub>4</sub>I after accounting for their common variation with burn size. Mean LO rose during the first week in nonsurvivors and remained elevated until death; their reduced FT<sub>4</sub>I apparently preceded the reduction in mental status. The relationship of nonsurvival to FT<sub>4</sub>I and LO did not depend upon administration of iodine, dopamine, corticosteroids, narcotic analgesics, or tranquilizers. Thus, we think it is reasonable at least to consider an alternate hypothesis for our results, in which persistently reduced circulating T<sub>4</sub> might have contributed to defective central nervous function and, possibly, to mortality. The measurement of obtundation is nonspecific, and further studies are necessary to test directly whether a deficiency of T<sub>4</sub> available to the brain in fatal NTI can contribute to either a pathophysiologically significant local deficit of hormone in that tissue or death.

### Acknowledgments

We wish to thank James Bullard, Len Seraile, and Sandy H. Coggins for technical and editorial assistance.

### References

1. Popp MB, Srivastava LS, Knowles Jr HC, MacMillan BG 1977 Anterior pituitary function in thermally injured male children and young adults. *Surg Gynecol Obstet* 145:517
2. Becker RA, Wilmore DW, Goodwin Jr CW, Zitzka CA, Wartofsky L, Burman KD, Mason AD, Pruitt BA 1980 Free T<sub>4</sub>, free T<sub>3</sub>, and reverse T<sub>3</sub> in critically ill, thermally injured patients. *J Trauma* 20:713
3. Becker RA, Vaughan GM, Ziegler MG, Seraile LG, Goldfarb IW, Mansour EH, McManus WF, Pruitt Jr BA, Mason Jr AD 1982 Hypermetabolic low triiodothyronine syndrome of burn injury. *Crit Care Med* 10:870
4. Wartofsky L, Burman KD 1982 Alterations in thyroid function in patients with systemic illness: the "euthyroid sick syndrome." *Endocr Rev* 3:164
5. Chopra IJ, Hershmann JM, Pardridge WM, Nicoloff JT 1983 UCLA Conference: thyroid function in nonthyroidal illnesses. *Ann Intern Med* 98:946
6. Slag MF, Morley JE, Elson MK, Crowson TW, Nuttall FQ, Shafer RB 1981 Hypothyroxinemia in critically ill patients as a predictor of high mortality. *JAMA* 245:43
7. Kaptein EM, Weiner JM, Robinson WJ, Wheeler WS, Nicoloff JT 1982 Relationship of altered thyroid hormone indices to survival in nonthyroidal illnesses. *Clin Endocrinol (Oxf)* 16:565
8. Calvano SE, Chiao J, Reaves LE, Antonacci AC, Shires T 1984 Changes in free and total levels of plasma cortisol and thyroxine following thermal injury in man. *J Burn Care Res* 5:143
9. Rudman D, Fleischer AS, Kutner MH, Raggio JF 1977 Suprahypophyseal hypogonadism and hypothyroidism during prolonged coma after head injury. *J Clin Endocrinol Metab* 45:747
10. Hill MA 1979 Annotated Computer Output for Regression Analysis. Health Sciences Computing Facility. Technical report no. 48, University of California, Los Angeles
11. Smeds S, Kagedal B, Lieden G, Liljedahl S-O 1981 Thyroid function after thermal trauma. *Scand J Plast Reconstr Surg* 15:141
12. Slag MF, Morley JE, Elson MK, Labrosse KR, Crowson TW, Nuttall FQ, Shafer RB 1981 Free thyroxine levels in critically ill patients. *JAMA* 246:2702
13. Melmed S, Geola FL, Reed AW, Pekary AE, Park J, Hershman JM 1982 A comparison of methods for assessing thyroid function in nonthyroidal illness. *J Clin Endocrinol Metab* 54:300
14. Kaptein EM, Robinson WJ, Grieb DA, Nicoloff JT 1982 Peripheral serum thyroxine, triiodothyronine and reverse triiodothyronine kinetics in the low thyroxine state of acute nonthyroidal illness. *J Clin Invest* 69:526
15. Chopra IJ, Huang T-S, Hurd RE, Beredo A, Solomon DH 1984 A competitive ligand binding assay for measurement of thyroid hormone-binding inhibitor in serum and tissues. *J Clin Endocrinol Metab* 58:619
16. Woebber KA, Maddux BA 1981 Thyroid hormone binding in nonthyroidal illness. *Metabolism* 30:412
17. Oppenheimer JH, Schwartz HL, Mariash CN, Kaiser FE 1982 Evidence for a factor in the sera of patients with nonthyroidal disease which inhibits iodothyronine binding by solid matrices, serum proteins, and rat hepatocytes. *J Clin Endocrinol Metab* 54:751
18. Eisenberg D, Silberman H, Ryan J, Shafer R, Weiss J, Kaptein EM, Nicoloff JT, Spencer CA 1980 Prognostic significance of thyroid function tests in critically ill patients. *Surg Forum* 31:211



NTIS	Dist	By	Availability Codes
DTIC TAB	Avail and/or	Distribution/	
Unannounced	Special		
Justification			

9-1 21